

AD-A155 583

ANTISTARSHAPEDNESS DISPERSIVENESS AND MIXTURES(U)
PENNSYLVANIA STATE UNIV UNIVERSITY PARK DEPT OF
STATISTICS J LYNCH 30 APR 85 ARO-20063.3-MA

1/1


UNCLASSIFIED

DAG29-84-K-0007

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963-A

Q

AD-A155 583

Antistarshapedness, Dispersiveness and Mixtures


by

James Lynch

April, 1985

Department of Statistics
The Pennsylvania State University

USARO Technical Report



This research was supported in part by the U.S. Army Research Office Grant No. DAAG29-84K-0007.

Key Words: Mixture, quantile plots, antistarshaped, dispersive, log concave.

AMS 1980 Subject Classification Numbers: Primary 62E10
Secondary 62N05

Approved for public release; distribution unlimited.

The view, opinions, and/or findings contained in this report are those of the author and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.

DTIC FILE COPY

6 6 0 5 8

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER ACO 20063.3-MA	2. GOVT ACCESSION NO. N/A	3. RECIPIENT'S CATALOG NUMBER N/A
4. TITLE (and Subtitle) Antistarshapedness, Dispersiveness and Mixtures		5. TYPE OF REPORT & PERIOD COVERED Technical
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) James Lynch		8. CONTRACT OR GRANT NUMBER(s) DAAG29-84K-0007
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office Post Office Box 12211 Research Triangle Park NC 27709		12. REPORT DATE April 30, 1985
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 7
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) NA		
18. SUPPLEMENTARY NOTES The view, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Mixture, quantile plots, antistarshaped, dispersive, log concave.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Necessary and sufficient conditions are given for certain classes of distributions to be closed under mixtures. The relevant classes are defined in terms of antistarshapedness or dispersiveness. One of the main results characterizes distributions with log concave densities.		

Abstract

Necessary and sufficient conditions are given for certain classes of distributions to be closed under mixtures. The relevant classes are defined in terms of antistarshapedness or dispersiveness. One of the main results characterizes distributions with log concave densities.



A/

1. Introduction.

A basic tool in the analysis of univariate survival data is the quantile plot, $(G^{-1}(t), F_n^{-1}(t))$, where G is a fixed distribution of interest and F_n^{-1} is the empirical quantile function. For large sample sizes $F_n \approx F$, where F is the distribution of the observations. So the behavior of the empirical quantile plot should be similar to $(G^{-1}(t), F^{-1}(t))$ which is essentially the same as $(t, G^{-1}F(t))$. The study of this behavior leads quite naturally to classes of distributions $\{F: G^{-1}F \text{ has property } P\}$. Such classes were first studied by Van Zwet (1964) when P was the property of convexity.

Certain well-known classes fall within this framework. When G is exponential the IFR (increasing failure rate), IFRA (increasing failure rate average), DFR and DFRA are the classes where property P is, respectively, convexity, starshapedness, concavity and antistarshapedness. The latter two classes are known to be closed under mixtures (Barlow and Proschan, 1975, Section 4.4). For the arbitrary case necessary and sufficient conditions are given in Leon and Lynch (1983).

Corresponding to each lifetime random variable (r.v.) X with distribution F is a r.v. $\hat{X} = \log X$ with distribution $\hat{F}(x) = F(e^x)$, $-\infty < x < \infty$. A distribution \hat{F} is said to be more dispersed than \hat{G} , written $\hat{G} \stackrel{d}{<} \hat{F}$, if

$$(1.1) \quad \hat{G}^{-1}(\beta) - \hat{G}^{-1}(\alpha) \leq \hat{F}^{-1}(\beta) - \hat{F}^{-1}(\alpha) \quad \alpha < \beta$$

This notion of dispersion is equivalent to $G^{-1}F$ being antistarshaped, i.e.,

$$(1.1') \quad \theta G^{-1}F(x) \leq G^{-1}F(\theta x) \quad 0 < \theta < 1, x > 0,$$

written $F \overset{a}{<} G$, and has been studied by a number of authors - Doksum (1969), Saunders and Moran (1978), Bickel and Lehmann (1979), Lewis and Thompson (1981) and Shaked (1982). Of particular note is Lewis and Thompson's result that the preservation of the dispersive ordering characterizes distributions with log concave densities.

In Section 2, we use the Lewis and Thompson characterization along with results in Leon and Lynch (1983) and the correspondence between (1.1) and (1.1') to find necessary and sufficient conditions on G for classes of distributions antistarshaped with respect to G to be closed under mixtures. Then using this same correspondence we obtain necessary and sufficient conditions for classes more dispersed than \hat{G} to be closed under mixtures, one of which is that \hat{G} have a log concave density.

2. The antistarshaped ordering and dispersiveness. In this section, the relationship between the antistarshaped ordering and the dispersive ordering is exploited to prove mixture results. The following definitions and lemma are needed. Throughout it is assumed that the support of the life distribution G is an interval $[0, c]$ if $c < \infty$ or $[0, \infty)$ if $c = \infty$.

Definition. A distribution \hat{G} is said to be two point dispersive if $\hat{G} \overset{d}{<} \hat{G} * \hat{F}$ whenever \hat{F} is a discrete distribution with only two mass points and where $*$ denotes the convolution operation. The distribution \hat{G} is said to be dispersive if $\hat{G} * \hat{F} \overset{d}{<} \hat{G} * \hat{H}$ whenever $\hat{F} \overset{d}{<} \hat{H}$.

Lemma 2.1. The following are equivalent:

- (1) \hat{G} is two point dispersive,

(ii) \hat{G} is dispersive, and

(iii) \hat{G} is absolutely continuous with a log concave density.

Proof (ii) \leftrightarrow (iii). This is in Lewis and Thompson (1981).

(ii) \rightarrow (i). This is obvious since it is easy to see that any distribution is more dispersed than a degenerate one.

(i) \rightarrow (iii). If \hat{G} is twice differentiable the argument in the proof of the only if part of Lewis and Thompson (1981) shows that \hat{G} has a log concave density. The proof then follows as in the last paragraph of Lynch, Mimmack and Proschan (1983). \square

Definition. The hazard transform for a life distribution G is given by

$$T_p(u, v) = G^{-1}(pG(u) + \bar{p}G(v))$$

where $0 < p < 1$, $\bar{p} = 1 - p$, and $0 \leq u, v < \infty$.

The following notation is needed. For two life distributions we denote the scale mixture $\int G(\lambda^{-1}x) dF(\lambda)$ by G_F .

Let

$$A_G = \{F: F \overset{a}{<} G\} \text{ and } \mathcal{D}_G^{\hat{}} = \{\hat{F}: \hat{G} \overset{d}{<} \hat{F}\}.$$

Thus $\mathcal{D}_G^{\hat{}} = \{\hat{F}: F \in A_G\}$ by the equivalence of (1.1) and (1.1').

The main results are given in the next two theorems.

Theorem 2.2. The following are equivalent:

- (i) A_G is closed under mixtures,
- (ii) $T_p(u, v)$ is antistarshaped for each p ,
- (iii) \hat{G} is dispersive (and so, \hat{G} has a log concave density), and

(iv) \mathcal{D}_G^\wedge is closed under mixtures.

Proof (i) \leftrightarrow (ii). This is just a restatement of Theorem 2.4(a) of Leon and Lynch (1983).

(ii) \leftrightarrow (iii). Because of Lemma 2.1, it suffices to show that (ii) is equivalent to \hat{G} being two point dispersive. Let $\hat{F} = p\delta_x + \bar{p}\delta_y$ where $\delta_x(z) = 0$ if $x \neq z$ and $= 1$ if $x = z$. Let $u = e^x$ and $v = e^y$. Then $F = p\delta_u + \bar{p}\delta_v$. Thus,

$$\begin{aligned} \hat{G} &\stackrel{d}{<} \hat{G} * \hat{F} \stackrel{a}{\text{iff}} G_F < G \\ &\text{iff } G^{-1}G_F(\theta z) \geq G^{-1}G_F(z) \text{ for } \theta \in (0,1) \\ &\text{iff } T_p(uz, uz) \text{ is antistarshaped} \\ &\text{iff } T_p(u, v) \text{ is antistarshaped.} \end{aligned}$$

(i) \leftrightarrow (iv). Ignoring certain measure theory details which are easily resolved and left to the reader, for a measure μ on A_G , let $\hat{\mu} = \mu I^{-1}$ where $I: A_G \rightarrow \mathcal{D}_G^\wedge$ is the 1-1 mapping $I(F) = \hat{F}$. Then,

$$\hat{G} \stackrel{d}{<} \int H d\hat{\mu} \stackrel{a}{\text{iff}} \int H d\mu < G.$$

Thus A_G is closed under mixtures if and only if \mathcal{D}_G is closed under mixtures. \square

As an immediate consequence of Lemma 2.1 and the equivalence of (1.1) and (1.1') we have

Theorem 2.3. $G_F \stackrel{a}{<} G_H$ whenever $F \stackrel{a}{<} H$ if and only if \hat{G} is dispersive.

Corollary 2.4. Let \hat{G} be dispersive. If $F \stackrel{a}{<} H$, then $G_F - G_H$ has at most one sign change and if one occurs it is from $-$ to $+$. If, in addition, $\int \lambda dF(\lambda) = \int \lambda dH(\lambda)$, then $G_F - G_H$ has exactly one sign change.

Proof. The proof is the same as for the DFRA case, i.e., G is the exponential distribution which is in Barlow and Proschan (1975). It is reproduced here for completeness.

Since $G_F \overset{a}{<} G_H$ by Theorem 2.3, $G_F - G_H$ has at most one sign change and it is from - to + if one occurs. If $\int \lambda dF(\lambda) = \int \lambda dH(\lambda)$, then $\int x dG_F(x) = \int x dG_H(x)$. Thus, $G_F - G_H$ must have at least one sign change. \square

Remark. Corollary 2.4 should be contrasted with Theorem 4 of Shaked (1980) where the ordering on the mixing distributions and the mixed distributions is the dilation ordering and with Theorem 1 of the same paper.

References

- Barlow, R. E. and Proschan, F. (1975). Statistical Theory of Reliability and Life Testing: Probability Models. Holt Rinehart and Winston, Inc., New York.
- Bickel, P. J. and Lehmann, E. L. (1979). Descriptive statistics for nonparametric models IV: spread. Contributions to Statistics - Jaroslav Hajek Memorial Volume, 33-40.
- Doksum, K. (1969). Starshaped transformations and the power of rank tests. Ann. Math. Statist. 40, 1167-1176.
- Leon, R. and Lynch, J. (1983). Preservation of life distribution classes under reliability operations. Math of OR 8, 159-169.
- Lewis, T. and Thompson, J. W. (1981). Dispersive distributions, and the connection between dispersivity and strong unimodality. J. Appl. Prob. 18, 76-90.
- Lynch, J., Mimmack, G. and Proschan, F. (1983). Dispersive ordering results. Adv. Appl. Prob. 15, 889-891.
- Saunders, I. W. and Moran, P. A. P. (1978). On the quantiles of the gamma and F distributions. J. Appl. Prob. 15, 426-432.
- Shaked, M. (1980). On mixtures from exponential families. J. R. Statist. Soc. B 42, 192-198.

Shaked, M. (1982). Dispersive ordering of distributions. J. Appl. Prob.
19, 310-320.

Van Zwet, W. R. (1964). Convex Transformations of Random Variables.
Amsterdam Mathematical Centre.

END

FILMED

8-85

DTIC